

An Observation of the Exoproducts in Media of Uropathogenic *E. coli*: Finding a Prospective Biological Mechanism for Uroepithelial Intracellular Colonization

Brittany N. Carver

BS Forensic Science; BS Biology-Pre Medical

Faculty Mentor: Donna L. Rhoads

Abstract

In this study, a procedure was developed to search for proteins secreted by UPEC that could be involved with urinary tract infections. UPEC strain CFT073 was inoculated in urine and incubated at body temperature for a total of 48 hours to imitate natural conditions in the human body. The bacterial cells were removed through centrifugation, and possible secreted proteins were isolated, quantified and separated using various protein analysis kits and SDS-PAGE. The results of the CFT073 strain were compared to a laboratory strain, MG1655 (K-12) and sterile urine. The data shows that there were no recognizable secreted proteins present in the UPEC urine cultures. This could indicate that secreted proteins larger than 5 kiloDaltons (kDa) are not involved in UTIs or that uroepithelial cell tissue is required for UPEC to secrete proteins.

Introduction

Women are more than 50% likely to contract a urinary tract infection (UTI) at least once in their lifetime.¹ As antibiotic resistance increases, the current methods of UTI treatment using ciprofloxacin and other antibiotics are becoming outdated, leaving patients at risk for recurrent urinary tract infections. When a UTI is asymptomatic or not treated, the infection can move from the bladder to the kidneys via the ureters causing pyelonephritis. Chronic urinary tract infections are a risk factor for bladder cancer. A recent study, done by Kyle Richards from the University of Wisconsin-Madison, showed that bladder cancer patients who experienced a UTI before their diagnosis were often at a more advanced stage in their cancer¹¹. In 2012, there were almost 600,000 people in the United States speculated to have bladder cancer.⁴ Understanding the mechanism of UTIs will lead to more efficient treatments, lower frequencies of pyelonephritis, and a lowered risk of bladder cancer.

Escherichia coli is a gram negative, facultatively anaerobic bacillus bacterium that is commonly found in the gut flora of mammals. Certain strains of *E. coli*, known as Uropathogenic *E. coli* (UPEC), are responsible “for more than 80% of uncomplicated UTIs”.² These strains have many virulence factors such as P fimbriae, hemolysin, serum resistance, and the ability to form quiescent intracellular bacterial reservoirs (QIR).³ These characteristics allow the bacterial cells to intracellularly infect uroepithelial host cells, cause urinary tract infections, provide the bacteria some antibiotic resistance, and affect the epigenetics of the host. P fimbriae which are found on the surface of the UPEC cells are known to adhere to receptors on the surface of uroepithelial cells. This adherence allows for the host cell to become infected. Other factors such as extracellular proteins, microRNA, and electrolytes could also be involved in the biochemical mechanism of the infection.

This study focuses on the protein output of UPEC strain CFT073 in urine compared to the laboratory strain K-12 due to their possible involvement in UTIs and the epigenetic effects on uroepithelial tissue. Exoproteins secreted by CFT073 should be distinguishable from *E. coli* K-12 and normal urine proteins. Results would then suggest that any found proteins have the potential for being involved in the biochemical mechanism that takes place during UTIs.

Materials and Methods

Urine was collected clean catch/mid-stream from a single donor, negative for a UTI, and filter-sterilized. The two strains, CFT073 and K-12, were inoculated in small samples varying from 3-4 mL of urine for 24 hours at body temperature. A cell counting experiment was performed on varying dilutions of the urine cultures by measuring absorbance at 600 nm (OD 600) using a spectrophotometer.⁵ The number of cells/mL were determined and related to absorbance. Next the difference in growth rates between CFT073 and K-12 was established by growing the strains in nutrient broth and sterile urine over a 48-hour period.

For protein isolation and quantification, two 24 hour 3 mL urine cultures of CFT073 and K-12 strains were made and used to find the OD 600. One milliliter of each culture was aseptically transferred to 3 mL of urine and incubated at 37°C for 24 hours. A Urine Protein Isolation and Concentration Kit by ITSI Biosciences was used to isolate and concentrate the protein from the urine. Before the procedure was performed, the samples were centrifuged at 12,000 rpm for 5 minutes and the supernatant was removed and transferred to fresh tubes to remove the bacterial cells from the extracellular protein and media.⁶ The isolated protein was analyzed using a Total Protein Assay Kit by ITSI Biosciences to create a standard curve and determine that protein concentration of the samples were within the limits of the sensitivity of SDS-PAGE.⁷ SDS-PAGE R-250 can detect as little as 100 ng¹².



Figure 1. The image to the right depicts the electrophoresis of a precast gel loaded with samples. Lane 1 - Urine Trial 2, Lane 3 - CFT073 Trial 2, Lane 5 - protein standard, Lane 7 - CFT073 Trial 1, Lane 9 - K-12 Trial 1, Lane 11 - protein standard, Lane 13 - K-12 Trial 2, Lane 15 - Urine Trial 1

An Edvotek vertical gel apparatus was used for SDS-PAGE. A Working Loading Buffer was prepared and added to the isolated protein samples at a ratio of 1:1. The mixture was boiled for 10 minutes to denature the proteins. Sigma-Aldrich TruePage Precast gels and SDS Running Buffer from ITSI Biosciences were added to the gel apparatus. Then samples and standards from BioRad were loaded into the gels, and electrophoresis was performed. After electrophoresis the gels were removed from the apparatus and soaked in a fixing/destaining solution. The fixed gels were placed in a Coomassie Brilliant Blue R-250 Staining Solution which binds to proteins. The stained gels were then placed in fixing/destaining solution until bands appeared by washing any unbound blue dye. The finished gels were placed in a storage solution which consisted of 0.05% acetic acid and distilled water and stored at 4°C.

Results

The cell counting experiment revealed that there were about $3.70E+08$ cells/mL in the CFT073 urine culture and that there were about $1.23E+06$ cells/mL in the K-12 urine culture. According to these results, there are about 300 times more cells per mL in a CFT073 culture than a K-12 incubated for 24 hours at 37°C in 3mL. Optical density (OD) was used to represent the measurement of the bacterial cell concentration at 600 nm. The average OD 600 at 24 hours for the K-12 strain was 0.12 and the average OD 600 for the CFT073 strain was 0.23.

The growth curve experiments conducted with urine had inconclusive results. Originally, the strains were inoculated into 3 mL of urine, but the optical density readings were low, and longer incubation times could not be

conducted. **Figure 2** shows the growth results of the strains after inoculation into the 0.360 mL wells of a 96 well plate.




Figure 2. Growth Curve results for 12 samples of K-12 and CFT073 in nutrient broth and urine over 48 hours at 37°C with constant orbital shaking in a 96 well plate. *UNEG* - Sterile Urine, *NBNEG* - Sterile Nutrient Broth (NB), *NBCFT* - CFT073 in NB, *NBK12* - K-12 in NB, *UK12* - K-12 in Urine, *UCFT1* - CFT073 in Urine.

NBCFT and *NBK12* represent the positive controls and show the growth curve that is expected for the two strains. *UNEG* and *NBNEG* represent the negative controls. The sample *UK12* has a very small growth curve, and *UCFT1*'s curve resembles that of the pure urine sample.

Significant data for K-12 was not obtained, therefore a statistical comparison between K-12 and CFT073 could not be performed. The average total extracellular proteins found for CFT073 was 770. µg/mL, and the average for the pure urine was 738 µg/mL. **Figure 3** shows the extracellular protein data for CFT073 and includes values for the pure urine samples and K-12. By using statistical tests to compare experimental means, *t*, was calculated to be 0.276 and the critical value of *t* was 2.06. This means that the two values are statistically equal because *t* critical is greater than the experimental *t*.




Figure 3. Average Total Extracellular Proteins in 48 hour cultures of CFT073 and K-12. The Standard Plot was determined with the Total Protein Assay Kit. The “X” on the plot designates the average total protein found in sterile urine. This figure demonstrates that CFT073 does not always secrete proteins in urine. This does not include proteins under 5 kDa.

The SDS-PAGE experiments all yielded gels that looked similar to the gel shown in **Figure 4**.




Figure 4. Finished SDS-PAGE gel. The well contents are labeled just above the figure. T1 and T2 represent Trial 1 and Trial 2. The molecular weights in kDa of the Prestained SDS-PAGE Standards are to the right of the figure lined up to the corresponding band.

Discussion

Since both strains were inoculated in a similar fashion in the cell counting experiment, the result can suggest that CFT073 has a higher growth rate than K-12 in urine. However, the results in **Figure 2** were obtained using the Epoch Microplate Spectrophotometer. The test samples had very low values which resembled the negative controls. At 24 hours, the microplate results showed that K-12 had an OD 600 of nearly 0.25, and the CFT073 culture had an OD 600 that was close to the negative controls. This result was unexpected because CFT073 is known to cause urinary tract infections whereas K-12 is not and should not be growing better than the UPEC strain in these conditions. Many factors could have contributed to this inaccuracy: a difference in inoculation technique, the size of the wells or contamination in the blank urine samples. Throughout the rest of the experiment, the bacteria were grown in 3-4 mL of urine and gave consistent readings at OD 600 that suggested growth. Urine has significantly fewer nutrients than nutrient broth, so the smaller volume of 350 μ L in the 96 well plate may not have had enough nutrients for the bacteria to grow as in the larger volumes of urine.

The total amount of Extracellular Proteins in 48 hour 4 mL cultures of both strains are shown in **Figure 3** However, since the averages between CFT073 and the sterile urine samples were similar it cannot be concluded that there were extracellular protein secretions in the media

of CFT073. The results demonstrated by the gel in **Figure 4** corroborate with the findings from the protein quantification analysis. In this gel and all other gels during the experiment, there was no significant pattern of banding that would suggest that CFT073 or K-12 secretes proteins in urine in the 48 hour incubation period. It was expected that there would be background banding from the urine, because a random urine sample has a protein concentration ranging from 0 to 20 mg/dL.⁸ There are many different types of trace amounts of protein found in urine ranging from uroepithelial tissue and glands, blood, and seminal proteins from the reproductive tract in males.⁹ According to a study on protein secretion and pathogenic bacterial infection, “The secretion of proteins into the extracellular milieu appears to be essential for bacterial multiplication in infected host tissues, but it is not required for microbial growth under laboratory conditions.”¹⁰

Conclusion

In this experiment, there was no evidence to suggest that there are exoproteins larger than 5 kDa secreted by the UPEC strain CFT073 in urine. It is possible that these proteins were transient and exist only long enough to assist in the infection. Since there was no uroepithelial tissue to infect, transient proteins would have dissociated. The limits of the experiment could have affected the results. Proteins under 5 kDa were not analyzed by the Total Protein Assay Kit and were separated from the samples that were loaded into the gels. The SDS-PAGE protein standard did not go below 5 kDa, and the 20.6 and 7.1 kDa standards did not show up on the gels. This suggests that there could have been inconsistencies with the electrophoresis procedure. The sensitivity of the materials used for SDS-PAGE could have prevented small quantities of protein from being noticed. Many bands on the gel are too light to be able to compare the blank samples to the test samples. The presence of uroepithelial cells may be necessary for UPEC to secrete proteins. Another explanation for these results is that extracellular proteins may not be involved in the infection. Other factors such as microRNA could be used in triggering the UPEC to infect the host. There is also a possibility that the P fimbriae are the only mechanism in which UPEC adheres to and infects uroepithelial cells. This research suggests that future investigations should focus on proteins that are secreted by CFT073 when exposed to uroepithelial cells. Other avenues of study could be on immediately produced small peptides or factors other than secreted proteins such as microRNA and electrolytes.

References

1. Foxman B, and Schaeffer A. (2012). Urinary Tract Infections in Adults. *National Kidney and Urologic Diseases Information Clearinghouse, National Institutes of Health. No. 12-2097*. Retrieved from <http://kidney.niddk.nih.gov/KUDiseases/pubs/utiadult/>

2. Background on Uropathogenic *E. coli*. *Uropathogenic Escherichia coli*, *Mobley Research Laboratory*. Retrieved from <http://www.umich.edu/~hltmlab/research/coli/>
3. Tolg C, Nesrin S, Cortese R, et al. (2011). Uropathogenic *E. coli* infection provokes epigenetic downregulation of CDKN2A (p16INK4A) in uroepithelial cells. *Laboratory Investigation*. 91:825-836.
4. Cancer of the Urinary Bladder. *Surveillance, Epidemiology, and End Results Program*. Retrieved September 1, 2015, from <http://seer.cancer.gov/statfacts/html/urinb.html>
5. Reynolds, J. (2011). Counting Bacteria. Retrieved September 1, 2015, from http://delrio.dccd.edu/jreynolds/microbiology/2421/lab_manual/counts.pdf
6. Urine Protein Isolation and Concentration Using the ITSIPREP Total Protein Isolation-Urine Kit (ToPI-U). (2011). Retrieved September 2, 2015, from http://www.itsibio.com/documents/Validated BioLab Procedures/Kits_2011/K-0019 ToPI-U/K-0019-10 Protein Isolation from Urine.pdf
7. Total Protein Assay Using the ITSIPREP ToPA Kit. (2011). Retrieved September 2, 2015, from http://www.itsibio.com/documents/Validated BioLab Procedures/Kits_2011/K-0014 ToPA/K-0014-20 ToPA.pdf
8. Protein urine test: MedlinePlus Medical Encyclopedia. (2015, August 3). Retrieved September 1, 2015.
9. Grant, G., & Everall, P. (1957, November 1). The Proteins of Normal Urine. Retrieved September 2, 2015.
10. Lee, V., & Schneewind, O. (2001). Protein Secretin and the Pathogenesis of Bacterial Infections. Retrieved October 25, 2015.
11. News, N. (n.d.). Persistent UTI Symptoms May Signal Bladder Cancer. Retrieved October 25, 2015.
12. What is the minimum amount of protein that can be easily... (n.d.). Retrieved October 25, 2015.

Acknowledgements

We sincerely thank the University of New Haven for funding and the opportunity to participate in the Summer Undergraduate Research Fellowship. We would also like to thank Professor Donna Rhoads for advising our project, donors Frank and Pat Carrubba for helping to fund our project, the Department of Biology and Environmental Science for our ability to use the laboratories and equipment, and Bobby Rousseau for urine media donation and the contribution of the K-12 protein quantification data.



Brittany N. Carver is a senior double majoring in Forensic Science and Biology – Pre Medical at UNH. She worked as a lab assistant for Anatomy and Physiology and currently for Microbiology.

Her interests are in epigenetics and environmental conservation, and she is interested in continuing on to medical school to eventually become a Forensic Pathologist or Medical Examiner.